

Supporting Information

Three-Dimensional Au Microlattices as Positive Electrodes for Li-O₂ Batteries

Chen Xu,[†] Betar M. Gallant,[‡] Phillip U. Wunderlich,[§] Timm Lohmann,[§] Julia R. Greer,[†]

[†]Division of Engineering and Applied Science, California Institute of Technology,
Pasadena, CA 91125

[‡]Division of Chemistry and Chemical Engineering, California Institute of Technology,
Pasadena, CA 91125

[§]Research and Technology Center, Robert Bosch LLC, Palo Alto, CA 94304




Figure S1. Cyclic voltammogram of as-fabricated Au microlattice and SEM of the surface of the sample. (a) CV of hollow Au microlattice in 0.5 M H_2SO_4 with a scan rate of 50 mV s^{-1} ; (b) SEM image of rough polycrystalline Au surface obtained via constant current electrodeposition.




Figure S2. XRD patterns of pristine Au microlattice.




Figure S3. SEM image of the surface of a Au microlattice electrode discharged at 210 nA cm⁻²_{true}. No noticeable formation of “toroids”.




Figure S4. FTIR spectrum of a microlattice after the first discharge, taken in a N₂ glovebox. The grey zone indicates Li₂CO₃ peaks. The peaks at 1340 cm^{-1} , 1200 cm^{-1} , 1136 cm^{-1} and 1060 cm^{-1} denoted by “*” are attributed to residual LiTFSI. This sample is not washed prior to characterization. The peak LiTFSI locations are in good agreement with Gowda *et al*¹.




Figure S5. Raman spectra of a microlattices after the 1st charge, 1st discharge, and after 3 cycles ending with a 4th discharge. References obtained from commercially available powders from Sigma Aldrich.

The powder references exhibit peaks at Raman shifts of 790 cm^{-1} and 1090 cm^{-1} for Li_2O_2 and Li_2CO_3 , respectively, which agrees well with literature.²⁻⁷ The predominant product after the 1st discharge was Li_2O_2 , as indicated by a peak in the Raman data at 790 cm^{-1} . No Li_2CO_3 peaks were observed. The FTIR spectrum showed several peaks centered at 1400 cm^{-1} and 860 cm^{-1} , which suggests the presence of a small amount of Li_2CO_3 . This discrepancy between the FTIR and the Raman data may arise from the C–O bond being more infrared active than Raman active.^{8,9} The increase in Li_2CO_3 can be readily observed by the emergence of a peak at 1090 cm^{-1} in the Raman data, which was absent after the 1st discharge (Figure 5). This finding is supported by the IR data where the intensity of the Li_2CO_3 peaks increased, along with that of HCO_2Li and $\text{CH}_3\text{CO}_2\text{Li}$, with cycling.

An interesting feature observed in the Raman spectra showed a peak at 750 cm^{-1} , which does not coincide with the previously observed peak for Li_2O_2 at 790 cm^{-1} , nor does it align with expected side reaction products such as LiOH , HCO_2Li and $\text{CH}_3\text{CO}_2\text{Li}$. One possibility is that the 750 cm^{-1} peak belongs to Li_2O_2 where the O – O bond strength is different from that of the Li_2O_2 formed on the 1st cycle. Varying positions for the O-O stretch in Li_2O_2 formed during discharge have been reported in literature, with values between 745 cm^{-1} ¹⁰ and 808 cm^{-1} ^{10,11}. The origins of this peak shift remain elusive; several factors that can potentially affect the bonding strength have been proposed, for example the crystallinity of Li_2O_2 , which can lead to peak broadening or peak disappearance,^{12,13} or hydration of Li_2O_2 , which can lead to a blue shift of up to 70

cm^{-1} .^{5,14} It is likely that in this work this shift is caused by the local contamination on Li_2O_2 surfaces from LiOH or other species.

References

- (1) Gowda, S. R.; Brunet, A.; McCloskey, B. D. Implications of CO₂ Contamination in Rechargeable Nonaqueous Li –. **2013**, 2–5.
- (2) Ottakam Thotiyl, M. M.; Freunberger, S. a; Peng, Z.; Bruce, P. G. The Carbon Electrode in Nonaqueous Li-O₂ Cells. *J. Am. Chem. Soc.* **2013**, *135*, 494–500.
- (3) Peng, Z.; Freunberger, S.; Chen, Y.; Bruce, P. A Reversible and Higher-Rate Li-O₂ Battery. *Science (80-.)*. **2012**, *337*, 563–566.
- (4) Gunasekara, I.; Mukerjee, S.; Plichta, E. J.; Hendrickson, M. A.; Abraham, K. M. Microelectrode Diagnostics of Lithium-Air Batteries. *J. Electrochem. Soc.* **2014**, *161*, A381–A392.
- (5) Veith, G. M.; Nanda, J.; Delmau, L. H.; Dudney, N. J. Influence of Lithium Salts on the Discharge Chemistry of Li–Air Cells. *J. Phys. Chem. Lett.* **2012**, *3*, 1242–1247.
- (6) Zakharchenko, T. K.; Kozmenkova, A. Y.; Itkis, D. M.; Goodilin, E. A. Lithium Peroxide Crystal Clusters as a Natural Growth Feature of Discharge Products in Li-O₂ Cells. *Beilstein J. Nanotechnol.* **2013**, *4*, 758–762.
- (7) Zhai, D.; Wang, H.-H.; Lau, K. C.; Gao, J.; Redfern, P. C.; Kang, F.; Li, B.; Indacochea, E.; Das, U.; Sun, H.; *et al.* Raman Evidence for Late Stage Disproportionation in a Li-O₂ Battery. *J. Phys. Chem. Lett.* **2014**, 140722114526001.

- (8) Nakamoto, K. Infrared and Raman Spectra of Inorganic and Coordination Compounds: Part B: Applications in Coordination, Organometallic, and Bioinorganic Chemistry; 2008; pp. 1–408.
- (9) Siebert, F.; Hildebrandt, P. *Vibrational Spectroscopy in Life Science*; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, 2008.
- (10) Thapa, A. K.; Shin, T. H.; Ida, S.; Sumanasekera, G. U.; Sunkara, M. K.; Ishihara, T. Gold–Palladium Nanoparticles Supported by Mesoporous B-MnO₂ Air Electrode for Rechargeable Li-Air Battery. *J. Power Sources* **2012**, *220*, 211–216.
- (11) Lu, Y.-C.; Gallant, B. M.; Kwabi, D. G.; Harding, J. R.; Mitchell, R. R.; Whittingham, M. S.; Shao-Horn, Y. Lithium–oxygen Batteries: Bridging Mechanistic Understanding and Battery Performance. *Energy Environ. Sci.* **2013**, *6*, 750.
- (12) Xie, J.; Yao, X.; Madden, I. P.; Jiang, D.-E.; Chou, L.-Y.; Tsung, C.-K.; Wang, D. Selective Deposition of Ru Nanoparticles on TiSi₂ Nanonet and Its Utilization for Li₂O₂ Formation and Decomposition. *J. Am. Chem. Soc.* **2014**, *136*, 8903–8906.
- (13) Zhai, D.; Wang, H.-H.; Yang, J.; Lau, K. C.; Li, K.; Amine, K.; Curtiss, L. a. Disproportionation in Li-O₂ Batteries Based on a Large Surface Area Carbon Cathode. *J. Am. Chem. Soc.* **2013**, *135*, 15364–15372.
- (14) Eysel, H. H.; Thym, S. RAMAN Spectra of Peroxides. *Zeitschrift für Anorg. und Allg. Chemie* **1975**, *411*, 97–102.